कोविड—19 सर्वोच्च प्राथमिकता संख्या—852 / पॉच—5—2020

प्रेषक,

प्राणेश चन्द्र शुक्ल, उप सचिव,

उत्तर प्रदेश शासन।

सेवा में,

1- महानिदेशक, चिकित्सा एवं स्वास्थ्य सेवायें, उ०प्र० लखनऊ।

2- महानिदेशक, चिकित्सा शिक्षा एवं प्रशिक्षण, उ०प्र० लखनऊ।

चिकित्सा अनुभाग-5

लखनऊ; दिनांक: 13 अप्रैल, 2020

विषय:—कोविड—19 के रोगियों हेतु रैपिड एंटीबॉडी आधारित ब्लड टेस्ट के संबंध में भारत सरकार की एडवाइजरी।

महोदय,

भारतीय आयुर्विज्ञान अनुसंधान परिषद (आई०सी०एम०आर०) द्वारा Rapid Antibody based blood test for Covid-19 प्रारम्भ करने के संबंध में निर्गत एडवाइजरी की प्रति संलग्न करते हुए मुझे यह कहने का निदेश हुआ है कि प्रश्नगत प्रकरण में उक्त एडवाइजरी के अनुसार अग्रेतर आवश्यक कार्यवाही करने का कष्ट करें।

संलग्नक-यथोक्त।

भवदीय.

(प्राणेश चन्द्र शुक्ल)

उप सचिव।

संख्या-852 (1) / पांच-5-2020, तद्दिनांक

प्रतिलिपि निम्नलिखित को सूचनार्थ एवं आवश्यक कार्यवाही हेतु प्रेषित:-

- 1. प्रमुख सचिव / विशेष सचिव, चिकित्सा शिक्षा विभाग, उ०प्र० शासन,
- 2. निदेशक, संचारी रोग, उ०प्र०।
- 3. गार्ड फाइल।

संलग्नक-यथोक्त।

(वंद प्रकाश राय) अनु सचिव।

INDIAN COUNCIL OF MEDICAL RESEARCH

DEPARTMENT OF HEALTH RESEARCH

Advisory to start rapid antibody based blood test for COVID-19 (4 April 2020)

Strategy for areas reporting clusters (containment zone) and in large migration gatherings/evacuees centres

Cases of Influenza Like Illness (ILI) to be monitored in health facilities. Any surge in cases to be monitored and brought to the notice of Surveillance Officer/CMO for additional investigation.

As a matter of abundant precautions, all symptomatic ILI persons should be advised home quarantine for 14 days.

At facility level, symptomatic ILI individuals to be tested using rapid antibody tests.

- Antibody test negative:
 - If warranted, confirm by real-time RT-PCR using throat/nasal swab.
 - RT-PCR negative: Likely non-COVID-19 ILI
 - RT-PCR positive: Confirmed COVID-19 Case and action as per protocol to be initiated for isolation, treatment and contact tracing.


OR

- If real-time RT-PCR not done, home quarantine and repeat antibody testing after 10 days of the last rapid antibody test.
 - Antibody test negative: Likely non-COVID-19 ILI.
 - Antibody test positive: there is possibility of recent infection, quarantine for another 10 days.
- Antibody test positive: After clinical assessment, treatment in hospital or isolation as per protocol. Action as per protocol to be initiated for contact tracing.

If symptoms worsen, refer to designated COVID-19 hospitals. When home quarantine is not feasible, consider facility-based quarantine.

General Guidelines:

- Healthcare workers doing the rapid antibody test to use gloves, mask, and head covers.
- Healthcare workers collecting throat/nasal swab to follow standard national infection control guidelines.
- The rapid antibody tests approved by US-FDA/CE-IVD or non-CE-IVD validated by ICMR-NIV with marketing approval by DCGI be used.
- In order to ensure that all such cases are monitored and necessary action is initiated with respect to infectious disease management, details of all test results shall be uploaded in ICMR portal.
- All such organizations are duty bound to register themselves to ICMR portal and upload the data in real-time.
- Failure to do so, they will be held liable to action under Disaster Management Act, 2005.




STRATEGY FOR USE OF RAPID ANTIBODY BASED BLOOD TEST

(4 April, 2020)

Cluster (Containment zone) & Large migration gatherings/evacuees centres

Symptomatic ILI (Influenza-Like-Illness)
Cough, Cold, Low grade fever, Sore throat

Home quarantine for 14 days and \blacksquare



If symptoms worsen, refer to designated COVID-19 hospitals